



**Life
Course
Centre**

WORKING PAPER SERIES

No. 2023-02
February 2023

The Kids are Not Alright: Differential Trends in Mental Ill-Health in Australia

Ferdi Botha

Melbourne Institute: Applied Economic & Social Research, The University of Melbourne

Richard W. Morris

Central Clinical School, Faculty of Medicine and Health, University of Sydney
School of Psychology, Faculty of Science, University of Sydney

Peter Butterworth

Melbourne Institute: Applied Economic & Social Research, The University of Melbourne
National Centre for Epidemiology and Population Health, The Australian National University

Nick Glozier

Central Clinical School, Faculty of Medicine and Health, University of Sydney

The Australian Research Council Centre of Excellence
for Children and Families over the Life Course

Phone +61 7 3346 7477 Email lcc@uq.edu.au
lifecoursecentre.org.au



Australian Government
Australian Research Council



THE UNIVERSITY
OF QUEENSLAND
AUSTRALIA



THE UNIVERSITY OF
SYDNEY



THE UNIVERSITY OF
WESTERN
AUSTRALIA



THE UNIVERSITY OF
MELBOURNE

Non-Technical Summary

There have been clear deteriorations in the mental health of Australians since around 2010, with most of this worsening believed to have occurred among young adults under age 35. However, to date it remains unclear whether these adverse mental health trajectories are because of age itself, or whether it might be linked instead to when people were born rather than how old they are. Given this observed deterioration in mental health among Australians over the past decade, this study investigates to what extent this deterioration differs in people born in different decades. We therefore test for possible cohort differences in the mental health of Australians.

Using 20 years of data from the Household, Income and Labour Dynamics in Australia (HILDA) Survey, we find that the observed deterioration in mental health in the Australian population over time is most consistent with a cohort effect rather than a temporary age-effect. Notably, it is those individuals from the more recent cohorts, especially those born in the 1990s and to a lesser extent those born in the 1980s, who show the strongest trajectories of worsening mental health over time. Individuals in this cohort report worse mental health than earlier cohorts at the same ages. There is little evidence that mental health is worsening with age for people born prior to the 1980s. Because our model allowed us to predict future trajectories based on the current trajectory, we expect this decline in the most recent generations will continue as they age. The findings are similar for men and women, and the results are robust to alternative samples and measures used.

The findings from this study highlight that it is the poorer mental health of Millennials that is driving the apparent deterioration in population-level mental health. Understanding the context and changes in society that have differentially affected younger people may inform efforts to ameliorate this trend and prevent it continuing for emerging cohorts.

About the authors

Ferdi Botha is a Senior Research Fellow at the Melbourne Institute: Applied Economic & Social Research at the University of Melbourne, and a Research Fellow at the ARC Centre of Excellence for Children and Families over the Life Course. He is also a Co-Editor of Social Indicators Research. His research interests include subjective and financial well-being, economic sociology, mental health, and applied microeconometrics. His research has been published in international peer-reviewed journals such as Social Science & Medicine, Social Indicators Research, and Journal of Population Economics. Email: ferdi.botha@unimelb.edu.au

Richard W. Morris is a Senior Research Fellow with joint appointments in the School of Medicine and School of Psychology at the University of Sydney. He is an expert in computational cognitive models, and has received a NARSAD Young Investigator Award and NHMRC funding to conduct research on learning and decision-making in psychiatric illness. His research has been published in SSM–Population Health and BMJ Open. Email: richard.morris@sydney.edu.au

Peter Butterworth is a Professor at the Centre for Research on Ageing, Health and Wellbeing in the Research School of Population Health at The Australian National University. His research experience is in (psychiatric) epidemiology, psychology and biostatistics. His broad research interests include the social causes and social consequences of common mental disorders across the life course. His research has been published in journals such as Social Science & Medicine, Social Psychiatry and Psychiatric Epidemiology, and Journal of Affective Disorders. Email: peter.butterworth@anu.edu.au

Nick Glozier is a consultant psychiatrist specialising in epidemiology, trials, and health services research. His research interests are in public mental health, function, sleep, the workplace, stress, disability and discrimination in our changing psychosocial and technological environments. He is a Chief Investigator of the Life Course Centre at the University of Sydney and leader of the Life Course Centre's People Program. His research has been published in journals such as The Lancet Psychiatry, Social Science & Medicine, and Social Psychiatry and Psychiatric Epidemiology. Email: nick.glozier@sydney.edu.au

Acknowledgements

This paper uses unit record data from Household, Income and Labour Dynamics in Australia Survey [HILDA] conducted by the Australian Government Department of Social Services (DSS). The findings and views reported in this paper, however, are those of the authors and should not be attributed to the Australian Government, DSS, or any of DSS' contractors or partners. DOI: 10.26193/YP7MNU. This research was supported by the Australian Government through the Australian Research Council's Centre of Excellence for Children and Families over the Life Course (Project ID CE200100025).

DISCLAIMER: The content of this Working Paper does not necessarily reflect the views and opinions of the Life Course Centre. Responsibility for any information and views expressed in this Working Paper lies entirely with the author(s).

Abstract

Given the observed deterioration in mental health among Australians over the past decade, this study investigates to what extent this differs in people born in different decades – i.e., possible cohort differences in the mental health of Australians. Using 20 years of data from the Household, Income and Labour Dynamics in Australia (HILDA) Survey, we find strong evidence that cohort effects are driving the increase in population level mental ill-health. Deteriorating mental health is particularly pronounced among people born in the 1990s and seen to a lesser extent among the 1980s cohort. There is little evidence that mental health is worsening with age for people born prior to the 1980s. The findings from this study highlight that it is the poorer mental health of Millennials that is driving the apparent deterioration in population-level mental health. Understanding the context and changes in society that have differentially affected younger people may inform efforts to ameliorate this trend and prevent it continuing for emerging cohorts.

Keywords: Mental ill-health, cohort effects, trajectories, Australia

Suggested citation: Botha, F., Morris, Richard W., Butterworth, P. & Glozier, N. (2023). 'The kids are not alright: Differential trends in mental ill-health in Australia', Life Course Centre Working Paper Series, 2023-02. Institute for Social Science Research, The University of Queensland.

Introduction

There is recent evidence from many countries that population mental health has worsened over time, even prior to the COVID-19 pandemic. In Australia, this pattern is most strikingly illustrated by the increasing rates of reported mental and behavioural disorders in the regular National Health Survey series, increasing from 9.6% of Australians aged 15 years and over in 2001 to 20.1% in 2017/18 (Australian Bureau of Statistics, 2018) and increasing even further to 21.4% in 2020/21 (Australian Bureau of Statistics, 2022). Such evidence of worsening mental health is consistent with data showing the increasing use of both psychotropics and therapeutic services within populations (Peach et al., 2022). The worsening mental health over time is also shown in measures of psychological distress, including research using large longitudinal panel surveys such as the Household, Income and Labour Dynamics in Australia (HILDA) Survey, which showed a broad increase in overall rates of psychological distress (Kessler-10 scores) in Australia from 4.8% to 7.4% between 2007 to 2017 across the 18 to 64 age-ranges (Butterworth, Watson, & Wooden, 2020). Most other OECD countries have observed similar worsening in population levels of mental health, particularly among young people in the UK, USA, Netherlands, and Japan (Hidaka, 2012; Nishi, Susukida, Usuda, Mojtabai, & Yamanouchi, 2018; Ormel, Hollon, Kessler, Cuijpers, & Monroe, 2022; Twenge, Cooper, Joiner, Duffy, & Binau, 2019), though this pattern is not ubiquitous (e.g., not in Canada: Patten et al., 2016). This paper seeks to better understand the factors potentially driving this increase in mental ill-health among Australians over time, in relation to period, age and cohort effects.

The worsening of population mental health over time may be a *period effect* to the extent that it reflects a ubiquitous change experienced by all groups in the population at the same point in time, regardless of age. This could, for example, reflect a change in risks that affect everyone (e.g., climate change). Alternatively, recent widespread international disruptions, such as the global financial crisis (GFC) or the COVID-19 pandemic which both resulted in loss of economic opportunity, may have also broadly impacted on the mental health of all (Butterworth, Schurer, Trinh, Vera-Toscano, & Wooden, 2022; Glozier, Morris, & Schurer, 2022).

An overall worsening of population mental health over time may be a consequence of *age effects* in the context of changing population age structures, e.g., associated with population ageing (Baxter et al., 2014; Ferrari et al., 2013; Vos et al., 2016). Such 'age effects' reflect differences in rates of poor mental health tied to age but independent of the period and cohort. Comparing age groups over the population has revealed a U-shaped pattern in mental wellbeing in large cross-sectional surveys. These hedonic aspects of wellbeing (often measured using questions similar to those used to assess distress, but with a different valence) decline from young age groups (e.g., 18-20) to middle-age (50-55) before increasing to a peak at 70-75, although there are cultural and national differences (Steptoe, Deaton, & Stone, 2015; Stone, Schwartz, Broderick, & Deaton, 2010). In Australia, Burns, Butterworth, & Crisp (2020) evaluated age-related changes in the mental health of Australian adults using 17 years of HILDA data (2001-2017). Using the mental health inventory (MHI-5) derived from the SF36 they report only very small differences in mental health over age-groups, but an emerging downward trend for the youngest (18-24) and very oldest adults (75+) in 2017 data, which suggests an *inverted* U-shaped pattern may be emerging across the age range.

In contrast to age-related changes, which reflect consistent variation in mental health over the lifespan irrespective of time, mental health may also vary by birth cohort. *Cohort effects* refer to variance over time that is specific to individuals born in or around certain years (e.g., generational differences between "Millennials" and "Baby Boomers"). Cohort differences in mental health are more likely to reflect widespread societal changes in risk factors or vulnerability that differentially affect cohorts, such as the penetration of social media, as they are associated with the person-specific differences which persist over the age-range.

Because of the linear dependency between age-, cohort-, and period-effects (Bell, 2020; Fienberg & Mason, 1978; Kratz & Brüderl, 2021), disentangling age-related effects from cohort effects in mental health is fraught, but crucial if we are to identify the groups most at risk and potentially target effective prevention or early treatment approaches. There is no technical way to solve the dependency and identify the unique effect of each in a linear model (Fienberg & Mason, 1978; Holford, 1983; Luo, 2013). Adding covariates changes the model but not the identification problem. The only way to solve this issue is by fiat; that is by conceding some constraint whose appropriateness cannot be tested (Fienberg, 2013;

Fienberg & Mason, 1985; Mason & Fienberg, 1985), or by focusing on nonlinear effects and possibly some interactions (Bell, 2020).

In this paper, we aim to distinguish whether the widely observed rise in mental ill-health, an observed period effect, in Australia is due to variation with age, or differences between birth cohorts. To remove the linear dependency and identify any differences in trends between cohorts, we model mental health (MHI-5 scores) for each cohort as a nonlinear smooth function of age. Comparisons between smooths allow us to determine whether the average level of mental health differs between cohorts (controlling for age), and whether the *trajectory* of mental health (over age) differs between cohorts.

Methods

Data and study design

This analysis draws on 20 annual waves of longitudinal data from the Household Income and Labour Dynamics in Australia (HILDA) survey. The HILDA Survey is a nationally representative household panel (aside from those in very remote Australia and those in non-private dwellings) that commenced in 2001 with 13,969 participants within 7,682 households. The study design follows all original household members over time, includes people who join households in which an original household member resides, and included a top-up sample (adding an additional 2,153 households) in 2011. Attrition rates from the study are low by international standards, with the re-interview rate increasing from 87% in wave 2, to over 95% in wave 8 and subsequently.

At each wave, data is collected through a face-to-face interview (with option for a telephone survey) and a separate self-completion questionnaire (SCQ). Given the key measures in the current study are drawn from the SCQ, the current sample is limited to those who completed the SCQ in a given year. For this analysis, the birth cohort of each person was defined by the decade of birth year (1940s, 1950s, 1960s, 1970s, 1980s, 1990s). Thus, persons can only contribute to a single birth cohort, but can be observed multiple times across survey years/ages. Persons were excluded if they were born prior to 1940 or after 1999 due to inadequate sample sizes. Demographic details of the sample are provided in Table 1.

Mental Ill-Health Measure

The MHI-5 is a subscale of five items assessing positive and negative aspects of mental health from the SF-36 (Batterham, Sunderland, Slade, Calear, & Carragher, 2018; Cuijpers, Smits, Donker, Ten Have, & Graaf, 2009; Ware Jr, 2000; Ware Jr & Sherbourne, 1992). It is well-validated as a screening instrument or dichotomised to provide a proxy of common mental disorders in population research (Hoeymans et al., 2004; Rumpf et al., 2001). Respondents are asked to state how often they have experienced each of the following during the past four weeks:

1. “Been a nervous person”
2. “Felt so down in the dumps nothing could cheer you up”
3. “Felt calm and peaceful”
4. “Felt down”
5. “Been a happy person”

In accordance with the manual (Ware, Snow & Kosinski 2000), items were recoded so that higher scores indicated better mental health. Raw scores were summed across the items and then transformed to a 0-100 scale. A person-specific score was estimated in any year on which there were valid responses on three or more items, the average being calculated and applied to missing data.

In sensitivity analysis, we repeat the key analysis using the 10-item Kessler scale of psychological distress (K10) that has been included in every second wave of the HILDA Survey since 2007.

Analysis

We estimate penalized smooth trends for each cohort using restricted maximum likelihood (REML) in a generalized additive mixed modelling (GAMM) setting (Wood, 2004, 2006, 2011; Wood, Pya, & Säfken, 2016). This is an analogue to a linear multilevel model with varying intercepts and slopes among the cohorts, but here the slopes are allowed to “wobble”. The model includes a global smoothing term for the effect of age as well as cohort-specific terms,

so each cohort is allowed to have its own functional response, but the penalty ensures that functions too far from average are penalized.

Each smoother f_k is represented by a sum of p simpler, fixed basis functions. The basis functions (splines) were estimated by quadratically penalized likelihood maximization for automatic smoothness selection, with a starting value of $p = 9$:

$$y_{it} = \beta_k(\text{cohort}_i) + f(\text{age}_{it}) + f_{[k]}(\text{age}_{it}) + \zeta_i + \epsilon_{it}$$

$$\epsilon_i \sim N(0, \phi\sigma^2)$$

Where y_{it} is the continuous MHI-5 score for each person i over age t ; $\beta_{[k]}$ is the mean MHI-5 estimate for each $k = 1 \dots K$ birth cohort, after accounting for variations in trend over age; and $f_{[k]}$ are smooth functions for the trend in MHI-5 scores over age for each cohort.

The smooth trends were centered for identifiability reasons (Marra & Wood, 2012; Wood, 2013), however the resulting model estimation allowed two important comparisons: Firstly, the mean MHI-5 estimates ($\beta_{[k]}$) provided comparisons for the average difference in mental health between cohorts. However, interpreting these differences is difficult in the presence of trends over age in each cohort. For example, a mean difference could be due to a decreasing trend with age in one cohort or an increasing trend in the other cohort, rather than consistent differences in mental health over the age range. Thus, an important advantage provided by the current model are the centered $f_{[k]}$ smooth functions from which differences in trends between cohorts are directly estimated. The resulting difference smooths are also centered around zero and so mean differences in mental health are not accounted for by these smooths, but they will reveal whether mental health is changing with age in one cohort relative to the other, reference, cohort. The difference smooths also directly estimate the uncertainty around the difference, with confidence intervals that include the uncertainty about the mean difference as well as the centered smooth itself. This results in intervals with close to nominal (frequentist) coverage probabilities (Marra & Wood, 2012).

We did not compare cohorts more than a decade apart since there are few or no overlapping age groups observed, so we restricted ourselves to the five ($K - 1$) pairwise comparisons between each cohort and the next oldest cohort (i.e., the *reference* cohort).

To account for the person-level dependency when survey participants are measured more than once, we included a first-order autoregressive AR(1) term ϕ for the residuals based on the unique cross-wave ID for each person $i = 1 \dots I$, which is equivalent to including the person-level random intercept ζ_i nested within cohort. In sensitivity analyses we explored the influence of period effects, sex, and first interview, as well as comparisons with psychological distress. The results are presented in the Appendix.

Results

The demographics of each cohort are shown in Table 1. The characteristics associated with the latest observation from each person is presented.

Table 1. Demographics stratified by birth cohort

Characteristic	1940s, N = 2,791 ¹	1950s, N = 3,890 ¹	1960s, N = 4,564 ¹	1970s, N = 4,614 ¹	1980s, N = 6,133 ¹	1990s, N = 5,265 ¹
Female	1,417 (51%)	2,043 (53%)	2,385 (52%)	2,368 (51%)	3,118 (51%)	2,718 (52%)
Age (years)	72 (65, 75)	62 (55, 66)	53 (46, 56)	42 (34, 46)	31 (24, 35)	24 (21, 27)
MHI-5 score	80 (60, 88)	80 (64, 88)	76 (60, 84)	76 (60, 84)	72 (60, 84)	72 (56, 80)
Very high distress (K10 > 29)	82 (3.7%)	190 (6.3%)	243 (6.8%)	290 (8.4%)	407 (8.8%)	594 (13%)
Employment						
Employed	587 (21%)	1,993 (51%)	3,458 (76%)	3,646 (79%)	4,722 (77%)	3,779 (72%)
Not in labour force	2,189 (78%)	1,820 (47%)	928 (20%)	741 (16%)	1,000 (16%)	965 (18%)
Unemployed	15 (0.5%)	77 (2.0%)	178 (3.9%)	227 (4.9%)	411 (6.7%)	521 (9.9%)
Highest Ed.						
None	1,252 (45%)	1,201 (31%)	1,213 (27%)	808 (18%)	1,150 (19%)	1,140 (22%)
Highschool	1,044 (37%)	1,728 (44%)	2,165 (47%)	2,292 (50%)	3,147 (51%)	2,965 (56%)
Grad	490 (18%)	960 (25%)	1,184 (26%)	1,511 (33%)	1,832 (30%)	1,160 (22%)
Chronic illness	1,524 (55%)	1,518 (39%)	1,348 (30%)	999 (22%)	1,011 (16%)	945 (18%)
Relationship						
Married/De Facto	1,889 (68%)	2,750 (71%)	3,258 (71%)	3,197 (69%)	3,551 (58%)	2,346 (45%)
Separated/Divorced/Widowed	774 (28%)	866 (22%)	778 (17%)	427 (9.3%)	213 (3.5%)	37 (0.7%)
Single	127 (4.6%)	271 (7.0%)	528 (12%)	989 (21%)	2,367 (39%)	2,881 (55%)

¹n (%); Median (IQR). Total observations within each cohort: 1940s = 31,871; 1950s = 43,472; 1960s = 48,800; 1970s = 42,379; 1980s = 45,391; 1990s = 30,184.

Later cohorts in our sample are more likely to have poorer mental health (lower MHI-5 scores), higher distress, more likely to be single and unemployed, and less likely to be chronically ill or disabled.

The complete range of ages within each cohort, which includes every observation of each person in every year included in the final model, is shown in Table 2. This clearly demonstrates the overlap in age between adjacent cohorts.

Table 2. Age distribution by cohort

cohort	youngest	median	oldest	observations
1940s	52	66	80	31,871
1950s	42	56	70	43,472
1960s	32	46	60	48,800
1970s	22	37	50	42,379
1980s	15	27	40	45,391
1990s	15	21	30	30,184

Figure 1. Age and cohort effects on mental health over the past two decades


Figure 1 shows changes in MHI-5 scores in each survey year by age at time of survey (left panel), and the trends in each birth cohort as it ages (right panel), where the dotted line represents the average age effect ignoring cohort. Mental health is worse for younger age-groups in each survey year, but this age-related discrepancy is much greater in more recent

surveys (left panel), consistent with evidence of a cohort effect. The right panel shows that mental health is worse for more recent generations, where deviations from the dotted line indicate the presence of a cohort effect. In particular, *Millennials* (those born in the 1990s) have a lower score at the same age as earlier generations, and the later cohorts do not show the age-related increase seen in other earlier cohorts as they age. At age 30 the average MHI-5 score of those born in the 1990s is about 67 on the 0-100 scale, compared to about 72.5 and 74 for people born in the 1980s and 1970s respectively.

Some combinations of ages/years were not observed for all age-groups (left panel) or cohorts (right panel). For example, people born prior to 1940 were excluded and so the earliest year observed for the oldest age group (65-74) was 2006, and the left panel shows the trend line for that age group does not extend earlier than 2006. Likewise, the earliest age observed of people born in the 1940s was 52, and so the trend line for that cohort does not extend earlier than that age (right panel).

Some trend lines are flat (e.g., left panel, ages 65-74; right panel, 1960s cohort), which is a result of the penalised smoothing spline determining that no additional degrees of freedom are required to support curvature to explain the variance in that group over years/ages. The left panel suggests that the negative effect of time (survey year) on mental health gets smaller as age increases, and for those aged 65 and above there is an absence of a time trend. In the right panel, in contrast, the flat line for the 1960s cohort reflects that this is the middle point of cohorts when moving from worsening mental health with age for more recent cohorts and more distant cohorts showing improving mental health with increasing age.

Uncertainty is not quantified (e.g., confidence intervals) in these plots, but pairwise comparisons of the average difference between each cohort and the immediately prior cohort (reference cohort) is presented in Table 3. Moreover, Figure 2 presents the difference smooths for each pairwise comparison to statistically compare the trends over age between cohorts.


Table 3. Pairwise differences in average mental health between cohorts

contrast	conf.low	estimate	conf.high	p.value
90s vs. 80s	-4.094	-3.576	-3.059	0.000
80s vs. 70s	-2.381	-1.603	-0.824	0.000
70s vs. 60s	-1.594	-0.995	-0.395	0.001
60s vs. 50s	-1.294	-0.790	-0.287	0.002
50s vs. 40s	-1.326	-0.738	-0.150	0.014

There are significant pairwise differences between each cohort and its reference cohort ($p < .05$), indicating poorer mental health scores in the later cohort of each comparison. These results represent the mean differences in MHI-5 scores of each cohort, and as such interpreting these differences is difficult given the presence of age effects within each cohort. For example, the mean difference could be due to a decreasing trend with age in the later cohort, or an increasing trend in the earlier cohort, rather than differences in mental health over all ages. Pairwise comparisons of the smooth trends over age for each cohort are thus presented next.

Figure 2 shows the smooth trend estimates, along with 95% confidence intervals (which include the uncertainty about the overall mean as well as the centred smooth itself). In each row the earlier cohort is shown in the left column as the 'reference smooth', and the estimated difference between the reference cohort and the cohort born in the subsequent decade is shown in the right column as the 'difference smooth'. A significant difference in trend or slope is indicated by 95% confidence intervals which exclude zero (horizontal line) in opposite directions at each endpoint.

Figure 2. Centered estimates of cohort trajectories (left) and their differences to the subsequent cohort (right)



The trend in the centered difference smooths (right panels) reveals whether the *change* in MHI-5 scores, or slope, of the later cohort is significantly different from the slope of the earlier reference cohort (left panels) over the same age range, i.e., a cohort effect. For example, a significant negative slope in the right panel demonstrates MHI-5 scores are declining over time in the later cohort at a faster rate than the reference (earlier) cohort. However, a negative slope in the right panel does not by itself indicate whether average MHI-5 scores are deteriorating in that cohort as they age. Inspection of the reference cohort in the left panel is also necessary to determine whether the decline observed in the right panel represents a true deterioration in mental health. For example, the 1990s difference panel reveals the 1990s cohort's mental health trajectory is significantly declining with age relative to the 1980s cohort, and the 1980s smooth (reference in the right panel) is also significantly deteriorating relative to its own mean baseline. Together this represents evidence that MHI-5 scores in the 1990s cohort are *declining even faster* than the deteriorating mental health of the 1980s cohort. Compare this to the 1960s difference smooth in the right panel, where there is also a significant negative slope. Here the trend in the 1950s reference smooth (left panel) is positive, so the negative difference in the right panel is not due to changes in the 1960s cohort but rather improvement in mental health with age in the 1950s cohort. In general, no cohort shows a steeper decline relative to its reference than the 1990s cohort, and when cohorts prior to the 1990s cohort tend to decline relative to their earlier reference cohort (right panels) it is not due to a deterioration relative to their own mean baseline (left panels).

The statistical significance of the smooth differences indicates where the slope in differences between cohorts are non-zero (i.e., positive or negative). As such they reveal the presence of cohort-effects adjusted for age. Table 4 reports relevant p-values based on Nychka (1988). The p-values indicate that cohort effects exist between each of the recent adjacent cohorts, but the effect declines for earlier cohorts and is not evident between the earliest two cohorts examined (i.e., between the 1950s and 1940s cohorts).

Table 4. Approximate significance of smooth differences

term	k-index	edf	F.value	p.value
1990s difference	0.98	1.00	60.24	0.000
1980s difference	1.02	4.37	8.55	0.000
1970s difference	1.02	3.30	6.21	0.000
1960s difference	1.01	1.00	3.27	0.071
1950s difference	0.98	1.00	0.15	0.698

Sensitivity Tests

In addition to the main analysis, we conducted several sensitivity tests, the results of which are reported in the Appendix. First, we tested for the presence of any period effects. We added a smooth term for survey year to the model and recalculated the difference smooths between cohorts in a post-estimation procedure (instead of a direct estimation as done in the main analysis). Adding a smooth term for period did not substantially influence the difference smooths we report in Figure 2 (compare to Figure A2). Second, instead of using the MHI-5 as our measure of mental ill-health, we also modelled the Kessler-10 (K10) psychological distress scale which was collected in the HILDA Survey in alternate years from 2007 to 2019. The corresponding psychological distress trajectories for each cohort are shown in Figure A3, with higher K10 scores suggesting greater psychological distress. The trajectory patterns are consistent with (and essentially the mirror image of) those observed for the MHI-5 scale in Figure 1 (right panel), as psychological distress was higher for more recent cohorts than earlier cohorts at the same age. Third, we also modelled the prevalence of mental illness defined by an MHI-5 score below 52, and observed very similar trajectories, such that the prevalence of mental illness was higher in more recent than earlier cohorts adjusted for age (Figure A4). Fourth, we conducted a check for any gender differences in cohort effects of mental health (Figure A5). The results are very similar to those reported for the full sample in Figure 1. The intercepts for men and women are different, with men’s average mental health being better than women’s average mental health. However, men and women have similar mental health trajectory differences between cohorts. We also checked sensitivity of results to the addition of new respondents (via a top-up sample) that occurred in 2011 in the HILDA Survey (which

was performed to maintain representativeness of the survey). This was also around the same time that we start observing declines in mental health (Figure 1, left panel), and so we conducted an analysis excluding top-up sample members to confirm the declining mental health trajectories were not driven by the specific respondents in the top-up sample (Figure A6). The results did not change the cohort trajectories. Finally, we checked whether trajectories were influenced by social demand characteristics of the survey. Because people may be unwilling to provide poor mental health responses, especially in an unfamiliar survey or to a new interviewer, we excluded the first survey response for each individual and reconducted the analysis. The resulting pattern of cohort differences were somewhat muted due to the loss of variation but remained consistent with the main findings.

Discussion

Population mental health in Australia has been worsening over the past decade, even prior to the COVID-19 pandemic and its sequelae. This is especially the case for younger adults aged between 15 and 35. Although there has been constant debate about the possible drivers of these trajectories of worsening mental health (Patten et al., 2016; Jorm et al., 2017; Mulder et al., 2017), it is challenging to precisely identify the source of these patterns and the assumption often is that these are period effects.

Using 20 years of longitudinal data we modelled the changes to mental ill-health for each birth cohort in Australia in a flexible non-linear model. Our model allowed us to compare mental ill-health between cohorts, adjusted for age, and we find that the observed deterioration in mental health in the Australian population over time is most consistent with a cohort effect rather than a temporary age-effect. Importantly, it is those individuals from the more recent cohorts, especially the 1990s birth cohort (Millennials), who show the strongest trajectories of worsening mental health over time. Individuals in this cohort report worse mental health than earlier cohorts at the same ages. Because our model allowed us to predict future trajectories based on the current trajectory, we expect this decline in the most recent generations will continue as they age. Thus, the deterioration in mental health over time which has been reported in large cross-sectional surveys, likely reflects cohort-specific effects related to the experiences of young people born in the Millennial generation and, to

a lesser extent, those from the immediately prior cohort born in the 1980s. The findings are similar for men and women, and the results are robust to alternative samples and measures used. We think these recent trends are unlikely to spontaneously resolve without addressing the new or exaggerated risks that may be differentially affecting these recent cohorts.

While previous research has examined age effects in large population-based surveys, there are few examples that have specifically examined mental health. Many more studies have focused on subjective wellbeing, often measured by a single item life-satisfaction question and also known as *cognitive* wellbeing. This literature has found evidence of almost every plausible age-related trajectory in life satisfaction (Kratz & Brüderl, 2021), and shown that the pattern of results observed seems to depend on the statistical model adopted for analysis. In the current study, we employed a random intercept model and observed a U-shaped effect of age on mental health with middle-aged people having poorer mental health on average than younger or older people. Kratz and Brüderl (2021) showed that this pattern can be produced by a random intercept model which yields biased age effects due to endogenous selection of happier people with age. However, a sensitivity analysis that controlled for endogenous effects by mean centering MHI-5 scores within-person (and including the mean score as a predictor, Ligthart-Smith, 2016) also observed a U-shaped pattern in age effects, suggesting the results we report here are not necessarily biased by endogeneity. An important consideration is that the MHI-5 measures a distinct construct from cognitive wellbeing, (Diener et al., 2017); one which has a different response to major life events (Kettlewell et al., 2020).

We did not include major life events in our model because they can act as mediators that result from age and affect the response variable (i.e., MHI-5 scores) (Kettlewell et al., 2020). We also did not include other potential mediators of the effect of age on mental ill-health such as health status, relationship status, employment status, household income or region. As such our results should be considered a description of the *total* effect of age on mental health, rather than providing a causal explanation of the individual drivers of such trajectories. Our aim here was to describe the cohort-related differences rather than explain them. Likewise, our aim was not to build a prediction model to extrapolate beyond the range of

data, and instead we prefer to note the expansion of the appropriately adjusted 95% confidence intervals when estimating future observations for any particular cohort.

This study provides a starting point for more in-depth analysis, and we hope it will encourage other researchers to more closely examine the changes that have happened in mental ill-health in Australia in the last decade. This is apparent from the trends depicted in the left panel of Figure 1, showing the divergence in mental health beginning roughly at the same time the 1990s cohort would have entered the survey for the first time. Future research should aim to identify and build understanding of the causes of these patterns, such as whether later cohorts are less resilient to similar risk factors experienced by earlier cohorts or whether they experience more and/or a greater severity of risks for mental ill-health. Such evidence is critical if the deteriorating pattern of mental health is to be arrested or shifted.


Appendix

Period effects

Period effects refer to variance over time that is common across all age groups and cohorts, due to population-wide events such as the Global Financial Crisis (GFC) in 2008 or the COVID-19 pandemic that started in 2020. We estimated the nonlinear effect of period over the complete set of survey years (2000 to 2020) as a smooth term in a model with cohort and (smooth) age effects.

The centered smooth effect of period (after accounting for nonlinear age and cohort effects) is shown in Figure A1.

Figure A1. Smoothed effect of period




The results show a slow decline in average MHI-5 scores from 2007 but which becomes more exaggerated from around 2017. However, we have already shown that this trajectory is not the same across all age-groups or cohorts (e.g., Figure 1).

After including a smooth term for period, we estimated the smooth trajectory of mental health for each cohort, and in a post-estimation procedure calculated the difference between cohort smooths in a pairwise fashion (Figure A2). In contrast to the results reported in Figure 2, these difference smooths are not directly estimated and so while each comparison includes overlapping ages, age is not matched exactly in each pairwise comparison (up to ± 5 years lag may be present). Nevertheless, the pattern of differences supports the same inferences

drawn from Figure 2 and demonstrates that period effects are not an influential presence in the cohort differences we report here.


Figure A2. Cohort trajectories (left) and their differences to the subsequent cohort (right)



Psychological distress

K10 scores (psychological distress: where higher scores indicate greater distress) were collected in alternate years from 2007 to 2019. The corresponding trajectories for each cohort are shown in Figure A3. Psychological distress was higher for later cohorts than earlier cohorts at the same age.


Figure A3. Psychological distress (K10 scores)



Risk of mental illness

While the MHI-5 is not a diagnostic instrument, it has good psychometric properties when identifying DSM-V disorders in a community sample (AUC 0.877, Batterham, Sunderland, Slade, Calear, & Carragher, 2018). We used a cut-off value of 52 to identify people at risk of mental illness and estimated the prevalence of mental illness for each age-group and cohort.

Figure A4. Age and cohort effects on prevalence of mental illness risk



The prevalence of risk for mental illness varied between age groups and cohorts in a similar pattern as psychological distress. Risk of mental illness was higher for later cohorts than earlier cohorts at the same age.

Other sensitivity checks


The difference between male and female MHI-5 trajectories is mostly in the intercept (overall mean level). Men and women have similar trajectory differences between cohorts (see Figure A5).

Figure A5. Interaction between sex and cohort




Excluding the Wave 11 top-up sample does not greatly impact the cohort trajectories (Figure A6).

Figure A6. Excluding top-up sample



Excluding the first survey response from each individual due to social demand characteristics of the interview process. Note the SF-36 is part of the SCQ so no interviewer is present, and we expect social demand characteristics to be low.

Figure A7. Excluding each respondent's first survey response



References

- Australian Bureau of Statistics. (2018). National Health Survey: First results. ABS. Retrieved from <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/national-health-survey-first-results/2017-18>
- Australian Bureau of Statistics. (2022). National Study of Mental Health and Wellbeing. ABS. Retrieved from <https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/latest-release>
- Batterham, P., Sunderland, M., Slade, T., Cleave, A., & Carragher, N. (2018). Assessing distress in the community: Psychometric properties and crosswalk comparison of eight measures of psychological distress. *Psychological Medicine*, 48(8), 1316–1324. <https://doi.org/10.1017/S0033291717002835>
- Baxter, A. J., Scott, K. M., Ferrari, A. J., Norman, R. E., Vos, T., & Whiteford, H. A. (2014). Challenging the myth of an “epidemic” of common mental disorders: Trends in the global prevalence of anxiety and depression between 1990 and 2010. *Depression and Anxiety*, 31(6), 506–516. <https://doi.org/10.1002/da.22230>
- Bell, A. (2020). Age Period Cohort analysis: A review of what we should and shouldn't do. *Annals of Human Biology*, 47(2), 208–217. <https://doi.org/10.1080/03014460.2019.1707872>
- Burns, R. A., Butterworth, P., & Crisp, D. A. (2020). Age, sex and period estimates of Australia's mental health over the last 17 years. *Australian & New Zealand Journal of Psychiatry*, 54(6), 602–608. <https://doi.org/10.1177/0004867419888289>
- Butterworth, P., Schurer, S., Trinh, T.-A., Vera-Toscano, E., & Wooden, M. (2022). Effect of lockdown on mental health in Australia: Evidence from a natural experiment analysing a longitudinal probability sample survey. *The Lancet Public Health*. [https://doi.org/10.1016/S2468-2667\(22\)00082-2](https://doi.org/10.1016/S2468-2667(22)00082-2)
- Butterworth, P., Watson, N., & Wooden, M. (2020). Trends in the prevalence of psychological distress over time: Comparing results from longitudinal and repeated cross-sectional surveys. *Frontiers in Psychiatry*, 1345. <https://doi.org/10.3389/fpsy.2020.595696>
- Cuijpers, P., Smits, N., Donker, T., Ten Have, M., & Graaf, R. de. (2009). Screening for mood and anxiety disorders with the five-item, the three-item, and the two-item mental health inventory. *Psychiatry Research*, 168(3), 250–255. <https://doi.org/10.1016/j.psychres.2008.05.012>
- Diener, E., Heintzelman, S. J., Kushlev, K., Tay, L., Wirtz, D., Lutes, L. D., & Oishi, S. (2017). Findings all psychologists should know from the new science on subjective well-being. *Canadian Psychology/Psychologie Canadienne*, 58(2), 87. Retrieved from <http://psycnet.apa.org/record/2016-48141-001>
- Ferrari, A. J., Charlson, F. J., Norman, R. E., Patten, S. B., Freedman, G., Murray, C. J., ... Whiteford, H. A. (2013). Burden of depressive disorders by country, sex, age, and year:

Findings from the global burden of disease study 2010. *PLoS Medicine*, 10(11), e1001547.
<https://doi.org/10.1371/journal.pmed.1001547>

Fienberg, S. E. (2013). Cohort analysis' unholy quest: A discussion. *Demography*, 50(6), 1981–1984. <https://doi.org/10.1007/s13524-013-0251-z>

Fienberg, S. E., & Mason, W. M. (1978). Identification and estimation of age-period-cohort models in the analysis of discrete archival data. *Sociological Methodology*, 10, 1–67.
<https://doi.org/10.2307/270764>

Fienberg, S. E., & Mason, W. M. (1985). Specification and implementation of age, period and cohort models. In *Cohort analysis in social research* (pp. 45–88). Springer.
https://doi.org/10.1007/978-1-4613-8536-3_3

Glozier, N., Morris, R., & Schurer, S. (2022). What happened to the predicted COVID-19-induced suicide epidemic, and why? *Australian & New Zealand Journal of Psychiatry*, 00048674221131500. <https://doi.org/10.1177/00048674221131500>

Hidaka, B. H. (2012). Depression as a disease of modernity: Explanations for increasing prevalence. *Journal of Affective Disorders*, 140(3), 205–214.
<https://doi.org/10.1016/j.jad.2011.12.036>

Hoeymans, N., Garsen, A.A., Westert, G.P., & Verhaak, P.F. (2004). Measuring mental health of the Dutch population: A comparison of the GHQ-12 and the MHI-5. *Health and Quality of Life Outcomes*, 2, 23. <https://doi.org/10.1186/1477-7525-2-23>

Holford, T. R. (1983). The estimation of age, period and cohort effects for vital rates. *Biometrics*, 311–324. <https://doi.org/10.2307/2531004>

Jorm, A.F., Patten, S.B., Brugha, T.S., & Mojtabai, R. (2017). Has increased provision of treatment reduced the prevalence of common mental disorders? Review of the evidence from four countries. *World Psychiatry*, 16(1), 90-99. <https://doi.org/10.1002/wps.20388>

Kettlewell, N., Morris, R. W., Ho, N., Cobb-Clark, D. A., Cripps, S., & Glozier, N. (2020). The differential impact of major life events on cognitive and affective wellbeing. *SSM-Population Health*, 10, 100533. <https://doi.org/10.1016/j.ssmph.2019.100533>

Kratz, F., & Brüderl, J. (2021). *The age trajectory of happiness*. LMU Munich, Mimeo.

Lighthart-Smith, A. (2016). *A comparison of linear mixed models that include time-varying covariates*. Australian Government Department of Education and Training.

Luo, L. (2013). Assessing validity and application scope of the intrinsic estimator approach to the age-period-cohort problem. *Demography*, 50(6), 1945–1967.
<https://doi.org/10.1007/s13524-013-0243-z>

Marra, G., & Wood, S. N. (2012). Coverage properties of confidence intervals for generalized additive model components. *Scandinavian Journal of Statistics*, 39(1), 53–74. <https://doi.org/10.1111/j.1467-9469.2011.00760.x>

Mason, W. M., & Fienberg, S. (1985). *Cohort analysis in social research: Beyond the identification problem*. Springer Science & Business Media. https://doi.org/10.1007/978-1-4613-8536-3_1

Mulder, R., Rucklidge, J., & Wilkinson, S. (2017). Why has increased provision of psychiatric treatment not reduced the prevalence of mental disorder? *Australian and New Zealand Journal of Psychiatry*, 51(12), 1176–1177. <https://doi.org/10.1007/0004867417727356>

Nishi, D., Susukida, R., Usuda, K., Mojtabai, R., & Yamanouchi, Y. (2018). Trends in the prevalence of psychological distress and the use of mental health services from 2007 to 2016 in Japan. *Journal of Affective Disorders*, 239, 208–213. <https://doi.org/10.1016/j.jad.2018.07.016>

Ormel, J., Hollon, S. D., Kessler, R. C., Cuijpers, P., & Monroe, S. M. (2022). More treatment but no less depression: The treatment-prevalence paradox. *Clinical Psychology Review*, 91, 102111. <https://doi.org/10.1016/j.cpr.2021.102111>

Patten, S. B., Williams, J. V., Lavorato, D. H., Bulloch, A. G., Wiens, K., & Wang, J. (2016). Why is major depression prevalence not changing? *Journal of Affective Disorders*, 190, 93–97. <https://doi.org/10.1016/j.jad.2015.09.002>

Peach, N., Barrett, E., Cobham, V., Ross, J., Perrin, S., Bendall, S., ... Mills, K. (2022). The mental health of adolescents and young people experiencing traumatic stress and problematic substance use. *Drug and Alcohol Review*, 41, 16–17. <https://doi.org/10.1111/dar.13537>

Rumpf, H.J., Meyer, C., Hapke, U., & John, U. (2001). Screening for mental health: validity of the MHI-5 using DSM-IV Axis I psychiatric disorders as gold standard. *Psychiatry Research*, 105, 243–53. [https://doi.org/10.1016/S0165-1781\(01\)00329-8](https://doi.org/10.1016/S0165-1781(01)00329-8)

Steptoe, A., Deaton, A., & Stone, A. A. (2015). Subjective wellbeing, health, and ageing. *The Lancet*, 385(9968), 640–648. [https://doi.org/10.1016/S0140-6736\(13\)61489-0](https://doi.org/10.1016/S0140-6736(13)61489-0)

Stone, A. A., Schwartz, J. E., Broderick, J. E., & Deaton, A. (2010). A snapshot of the age distribution of psychological well-being in the United States. *Proceedings of the National Academy of Sciences*, 107(22), 9985–9990. <https://doi.org/10.1073/pnas.1003744107>

Twenge, J. M., Cooper, A. B., Joiner, T. E., Duffy, M. E., & Binau, S. G. (2019). Age, period, and cohort trends in mood disorder indicators and suicide-related outcomes in a nationally representative dataset, 2005–2017. *Journal of Abnormal Psychology*, 128(3), 185. <https://doi.org/10.1037/ABN0000410>

Vos, T., Allen, C., Arora, M., Barber, R. M., Bhutta, Z. A., Brown, A., ... others. (2016). Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases

and injuries, 1990–2015: A systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*, 388(10053), 1545–1602. [https://doi.org/10.1016/S0140-6736\(16\)31678-6](https://doi.org/10.1016/S0140-6736(16)31678-6)

Ware J. E. (2000). SF-36 health survey update. *Spine*, 25(24), 3130–3139.

Ware, J. E., Snow, K. K., & Kosinski, M. (2000). SF-36 version 2 health survey: Manual and interpretation guide. *Linc RI Qual Metr Inc*.

Ware J. E., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Medical Care*, 473–483.

Wood, S. N. (2004). Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association*, 99(467), 673–686. <https://doi.org/10.1198/016214504000000980>

Wood, S. N. (2006). Low-rank scale-invariant tensor product smooths for generalized additive mixed models. *Biometrics*, 62(4), 1025–1036. <https://doi.org/10.1111/j.1541-0420.2006.00574.x>

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(1), 3–36. <https://doi.org/10.1111/j.1467-9868.2010.00749.x>

Wood, S. N. (2013). On p-values for smooth components of an extended generalized additive model. *Biometrika*, 100(1), 221–228. <https://doi.org/10.1093/biomet/ass048>

Wood, S. N., Pya, N., & Säfken, B. (2016). Smoothing parameter and model selection for general smooth models. *Journal of the American Statistical Association*, 111(516), 1548–1563. <https://doi.org/10.1080/01621459.2016.1180986>